



MONTCLAIR STATE
UNIVERSITY

Montclair State University
**Montclair State University Digital
Commons**

Sustainability Seminar Series

Sustainability Seminar Series, 2018

Jan 18th, 4:00 PM - 5:00 PM

CO2 Levels And Climate Change During Early Eocene Hyperthermals

Ying Cui
Dartmouth College

Follow this and additional works at: <https://digitalcommons.montclair.edu/sustainability-seminar>



Part of the [Environmental Sciences Commons](#)

Cui, Ying, "CO2 Levels And Climate Change During Early Eocene Hyperthermals" (2018). *Sustainability Seminar Series*. 10.

<https://digitalcommons.montclair.edu/sustainability-seminar/2018/spring2018/10>

This Open Access is brought to you for free and open access by the Conferences, Symposia and Events at Montclair State University Digital Commons. It has been accepted for inclusion in Sustainability Seminar Series by an authorized administrator of Montclair State University Digital Commons. For more information, please contact digitalcommons@montclair.edu.



MONTCLAIR STATE
UNIVERSITY

The MSU Sustainability Seminar Series Presents:
CO₂ levels and climate change during Early Eocene hyperthermals

WHEN: January 18 (Thursday), 4:00 pm

WHERE: CELS 120 lecture hall

Dr. Ying Cui
Dartmouth College



Dr. Cui is currently an Obering Postdoctoral Fellow from the Department of Earth Sciences at Dartmouth College. She obtained a PhD in Geosciences and Biogeochemistry at Penn State. Her research deals with carbon cycles during ancient hyperthermal events, such as the Paleocene-Eocene Thermal Maximum and the end-Permian mass extinction event. Her main research interest is in exploring the links between extinction, carbon emission and hyperthermals. She has published 12 peer-review articles related to this research topic, including *Nature Geoscience*, *Earth and Planetary Science Letters*, *Geochimica et Cosmochimica Acta* and *Earth-Science Reviews*. She is currently funded by National Science Foundation to study the atmospheric $p\text{CO}_2$ and Earth system climate sensitivity during the Cenozoic.

Early Eocene (56 to 33.9 million years ago) is punctuated by a series of rapid and extreme global warming events, known as the 'hyperthermals', triggered by massive release of carbon dioxide. Despite the carbon emission rate might have been ten times smaller than the current emission from burning fossil fuels and cement production, the early Eocene hyperthermals are considered one of the best analogues for CO₂ concentration projected into the future. The temperature records are well established from deep sea sediments during these hyperthermals, but the CO₂ levels in the atmosphere remain poorly known, partly due to the lack of a reliable CO₂ proxy.

Thanks to a newly developed C₃ land plant CO₂ proxy, we were able to calculate the upper limit of the CO₂ levels just before and during five early Eocene hyperthermals (PETM, H1, H2, I1 and I2). This proxy is based on the empirical relationship between carbon isotope fractionation and the ambient CO₂ levels: i.e. as CO₂ level rises, carbon isotope fractionation also increases. By comparing the carbon isotope excursion recorded in the ocean and on land, we were able to find that CO₂ levels just before each of the hyperthermals may have been less than 1000 parts per million, or ppm. CO₂ level as high as 2600 ppm might have occurred only briefly during the hyperthermals. This suggests that to maintain the early Eocene ice-free condition and global average surface temperature of ~25 °C, other greenhouse gases, such as methane, water vapor and nitrous oxide must have played more important role.